

The large— N behavior of the holographic models of QCD¹

Frédéric Jugeau

*Institute of High Energy Physics, Chinese Academy of Sciences,
P. O. Box 918(4), Beijing 100049, China.
Theoretical Physics Center for Science Facilities, Chinese Academy of Sciences,
Beijing 100049, China (jugeau@ihep.ac.cn).*

Abstract. The AdS/QCD dictionary is considered, checking the large— N behavior of $5d$ dual models of QCD as a guideline. Especially, a consistent chiral symmetry breaking function in the Hard Wall model is derived and the different forms of the field/operator correspondence are emphasized.

Keywords: holographic models of QCD, large— N 't Hooft limit, pion form factors in the chiral limit.

PACS: 11.15.Pg, 11.25.Tq, 11.30.Rd, 14.40.Aq.

INTRODUCTION

There is a new hope that the AdS/CFT correspondence provides us a new way to deal with the non-perturbative regime of QCD. Any attempt to establish the AdS/QCD duality starting from the tested AdS/CFT recipe [1] should then lead to a consistent large— N behavior for the $5d$ holographic models of QCD, the tree level approximation of the dual string theory corresponding indeed to the 't Hooft limit of the gauge theory. Within the *bottom-up* approach, the so-called Hard Wall model [2] consists in breaking the underlying conformal symmetry by truncating the holographic AdS_5 space-time such as $0 < z \leq z_m \simeq 1/\Lambda_{QCD}$. Moreover, one usually assumes that only a small number of QCD operators are relevant for describing the low-energy phenomenology of the lightest mesons. Among them, the scalar bulk field $v(z)$ dual to the quark condensate $\langle \bar{q}q \rangle$ accounts for the chiral symmetry breaking (in the sequel, we shall work with units in which the AdS_5 radius is put equal to 1):

$$v(z) = z(m_q + \bar{\sigma}z^2) \xrightarrow{z \rightarrow 0} m_q z . \quad (1)$$

The source field m_q is nothing else than the light quark mass while $\bar{\sigma}$ is rather systematically identified in the literature with the chiral condensate $-\langle \bar{q}q \rangle$. As pointed out in [3], such a naive identification leads to inconsistencies regarding the large— N behavior since $m_q \sim O(N^0)$ and $\langle \bar{q}q \rangle \sim O(N)$. The aim of this talk is thus to reexamine the AdS/QCD dictionary, checking the large— N behavior of the Hard Wall model as a guideline. Doing

¹ Talk given at the 2nd annual *France China Particle Physics Laboratory* (FCPPL) Workshop, March 21-24, 2009, Wuhan, China.

so, we shall be able to identify unambiguously the quark mass and the chiral condensate entering the expression of the so-called chiral symmetry breaking function $v(z)$ [4].

THE CHIRAL SYMMETRY BREAKING FUNCTION AND THE *AdS/QCD* DICTIONARY

The $5d$ holographic space-time is described by the following conformally flat metric ($M, N = 0, \dots, 4$ and $\mu, \nu = 0, \dots, 3$):

$$ds^2 = g_{MN}(z)dx^M dx^N = e^{2A(z)} \left(\eta_{\mu\nu} dx^\mu dx^\nu - dz^2 \right) \quad (2)$$

where $\eta_{\mu\nu} = \text{diag}(+1, -1, -1, -1)$ is the flat metric tensor of the $4d$ Minkowski boundary space, $A(z) = -\ln(z)$ is the warp factor and $0 < z \leq z_m$. The effective action able to reproduce the chiral properties of QCD is

$$S_{5d} = -\frac{1}{2\kappa^2} \int d^5x \sqrt{g} \left(\mathcal{R} + \Lambda \right) + \frac{1}{k} \int d^5x \sqrt{g} \text{Tr} \left\{ |DX|^2 - m_5^2 X^2 - \frac{1}{4g_5^2} (G_L^2 + G_R^2) \right\} \quad (3)$$

with $g = \det(g_{MN})$ the determinant of the metric tensor, $\kappa^2 \equiv 8\pi G_5$ the $5d$ Newton constant and where the AdS_5 Ricci scalar $\mathcal{R} = -\frac{5}{3}\Lambda = 20$ is, according to our convention (2), positive *i.e.* the cosmological constant Λ is negative. The overall parameter k has the dimension of a length while the $5d$ gauge coupling constant g_5^2 is dimensionless. The relevant QCD operators are the $SU(2)_L \times SU(2)_R$ left-handed and right-handed currents $j_{L\mu}^a = \bar{q}_L \gamma_\mu t^a q_L$ and $j_{R\mu}^a = \bar{q}_R \gamma_\mu t^a q_R$ and the quark operator $\bar{q}_R^i q_L^j$ (when $n_F = 2$, $t^a = \frac{\sigma^a}{2}$ are the Pauli matrices with $a = 1, 2, 3$ and $i, j = 1, 2$). The bulk field content of the $5d$ dual theory consists then of the left-handed and right-handed gauge fields $A_{LM}^a(x, z)$ and $A_{RM}^a(x, z)$ and of the bi-fundamental scalar field:

$$X^{ij}(x, z) = \left(\frac{v(z)}{2} + S(x, z) \right)^{ik} \left(e^{2i\pi^b(x, z)t^b} \right)^{kj} \quad (4)$$

which is massive $m_5^2 = (\Delta - p)(\Delta + p - 4) = -3$ with $\Delta = 3$ and $p = 0$ and involves the bulk field $S(x, z)$ dual to the scalar mesons, the dimensionless pseudo-scalar field $\pi(x, z)$ and the background field $v(z)$, written as $v^{ik}(z) = v(z)\delta^{ik}$ in the isospin limit. The linearized equations of motion for the different bulk fields can be derived straightforwardly from the quadratic part of the $5d$ effective action (3). In particular, the ($4d$ Fourier transforms of the) pseudo-scalar modes are not independent but satisfy two coupled differential equations:

$$\partial_z \left(\frac{1}{z} \partial_z \tilde{\phi}^a(q, z) \right) - \frac{g_5^2 v(z)^2}{z^3} \left(\tilde{\phi}^a(q, z) - \tilde{\pi}^a(q, z) \right) = 0 \quad , \quad (5)$$

$$q^2 \partial_z \tilde{\phi}^a(q, z) - \frac{g_5^2 v(z)^2}{z^2} \partial_z \tilde{\pi}^a(q, z) = 0 \quad (6)$$

where ϕ contributes to the longitudinal part of the axial-vector bulk field $A_M^a = (A_L^a - A_R^a)/2$, defined as $\partial\phi = A - A_\perp$.

According to the Gell-Mann-Oakes-Renner (GMOR) relation $m_\pi^2 f_\pi^2 = 2m_q \sigma + O(m_q^2)$, the magnitudes of the pion and quark masses are related to the size of the chiral condensate $\sigma \equiv -\langle \bar{q}q \rangle$. In the chiral limit, the massless pions are the pseudo-Goldstone bosons associated with the spontaneous breaking of the chiral symmetry. They acquire a mass provided that the chiral symmetry is explicitly broken. On the other hand, the chiral symmetry breaking mechanism is holographically described by the background field $v(z)$ which reads in the Hard Wall model $v(z) = z(\bar{m} + \bar{\sigma}z^2)$. The two parameters $\bar{m}$ and $\bar{\sigma}$, which do not have to be identified *a priori* with the quark mass and the chiral condensate accounts for the explicit and implicit chiral symmetry breaking respectively and, as such, should be naturally proportional, on the one hand, to m_q and, on the other hand, to σ . It is worth deriving shortly the GMOR relation. When $q^2 = m_\pi^2$, Eq.(6) reads

$$m_\pi^2 \partial_z \phi(z) = \frac{g_5^2 v(z)^2}{z^2} \partial_z \pi(z) . \quad (7)$$

The normalizable modes of the pions behave as $\phi(0) = 0$, $\partial_z \phi(z)|_{z_m} = 0$ and $\pi(0) = 0$ while its IR boundary condition is dictated by the chiral symmetry (the subscript χ refers to the chiral limit): since $\pi_\chi(z) = -1$ is solution of Eq.(6) when $q^2 = 0$, $\pi(z)$ must approach -1 at low energies. Then, at the leading order, Eq.(7) becomes:

$$\pi(z) = m_\pi^2 \int_0^z du f(u) \frac{1}{g_5^2 u} \partial_u \phi_\chi(u) \quad (8)$$

where the function $f(u) \equiv \frac{u^3}{v(u)^2}$ exhibits a peak located at $u_c = \sqrt{\frac{\bar{m}}{3\bar{\sigma}}}$. So, only small values of u give sizeable contributions to the integral provided that $\bar{m}$ is small in comparison with $\bar{\sigma}$, which is consistent with the fact that $\bar{m}$ and $\bar{\sigma}$ are related to m_q and σ respectively. The remaining function can then be evaluated at $u \simeq 0$ which yields the pion decay constant $f_\pi^2 = -\frac{1}{kg_5^2} \frac{1}{z} \partial_z \phi_\chi(z)|_{z=0}$. Finally, in order to recover the GMOR relation at leading order, one takes the large z limit which gives rise to the simple condition $\bar{m}\bar{\sigma} = km_q\sigma$ (9). It is then consistent to take $\bar{m} = m_q$ (10) and $\bar{\sigma} = k\sigma$ (11). On the other hand, since any holographic model must predict the same high energy physics near the UV brane (when $z \rightarrow 0$), the same value of k should be extracted within any dual model of QCD. The short distance *AdS/QCD* matching of the two-point correlation function of the scalar current gave, in the framework of the Soft Wall model, the value $k = \frac{16\pi^2}{N}$ [5]. The chiral symmetry breaking function takes then the following form:

$$v(z) = z \left(m_q + \frac{16\pi^2}{N} \sigma z^2 \right) \xrightarrow{z \rightarrow 0} m_q z . \quad (12)$$

Its large- N scaling is consistent since $m_q \sim O(N^0)$ and $\sigma \sim O(N)$ yield $v(z) \sim O(N^0)$.

Let us now clarify the *AdS/QCD* dictionary. A mere rescaling of the scalar bulk field $X(x, z)$ and of the $5d$ coupling constant :

$$\hat{X}(x, z) = \frac{a}{2} X(x, z) \quad \text{and} \quad \frac{1}{\hat{g}_5^2} = \frac{a^2}{4} \frac{1}{g_5^2} , \quad (13)$$

where the parameter $a = \sqrt{N}/2\pi$ has been introduced in [3], implies a renormalization of the $5d$ holographic action (3):

$$\frac{1}{k} \int d^5x \sqrt{g} \text{Tr} \mathcal{L}(X, g_5^2, A_L, A_R) \Rightarrow \int d^5x \sqrt{g} \text{Tr} \hat{\mathcal{L}}(\hat{X}, \hat{g}_5^2, A_L, A_R) . \quad (14)$$

The action, written in the first form, involves an overall factor $1/k$ related to the number of colors N . Because of the scalar bulk field and gauge coupling redefinition, this N -dependent factor is absorbed on the *r.h.s.* of (14). That implies modifications in the field/operator prescription, the chiral symmetry breaking function becoming indeed:

$$\hat{v}(z) = \frac{a}{2} m_q z + \frac{2}{a} \sigma z^3 \xrightarrow{z \rightarrow 0} \frac{a}{2} m_q z . \quad (15)$$

We have then $\bar{m} = \frac{a}{2} m_q$ and $\bar{\sigma} = \frac{2}{a} \sigma$ which fulfils the condition $\bar{m} \bar{\sigma} = m_q \sigma$, analogous to (9). Moreover, the large- N behavior has changed, being now $\hat{v}(z) \sim O(\sqrt{N})$. Besides, one usually chooses to renormalize the $5d$ gauge coupling such that $\hat{g}_5^2 = 12\pi^2/N$ instead of the gauge bulk fields $A_{L_M}^a$ and $A_{R_M}^a$ (or, equivalently, V_M^a and A_M^a) which has the advantage of letting unchanged the boundary conditions:

$$\tilde{V}(q^2, 0) = \tilde{A}(q^2, 0) = 1 \quad \text{and} \quad \partial_z \tilde{V}(q^2, z)|_{z=z_m} = \partial_z \tilde{A}(q^2, z)|_{z=z_m} = 0. \quad (16)$$

Also, the pseudo-scalar modes are not affected by the rescaling (13). As a result, all the $5d$ equations of motion and the boundary conditions, especially $\pi_\chi(z) = -1$, remain the same with g_5^2 and $v(z)$ replaced by $\hat{g}_5^2$ and $\hat{v}(z)$ such that the products $g_5^2 v(z)^2$ and $\hat{g}_5^2 \hat{v}(z)^2$, as in (5)-(6) for instance, are both N -independent in the chiral limit. As for the vector, axial-vector and pseudo-scalar decay constants, which read respectively

$$F_{V_n}^2 = \frac{1}{kg_5^2} \left(\frac{1}{z} \partial_z v_n(z) \right)^2 \Big|_{z=0} , \quad F_{A_n}^2 = \frac{1}{kg_5^2} \left(\frac{1}{z} \partial_z a_n(z) \right)^2 \Big|_{z=0} , \quad f_\pi^2 = -\frac{1}{kg_5^2} \frac{1}{z} \partial_z \phi_\chi(z) \Big|_{z=0} \quad (17)$$

with $v_n(z) = \sqrt{2} \frac{z}{z_m} \frac{J_1(m_{V_n} z)}{J_1(m_{V_n} z_m)}$ and $a_n(z)$ the vector and axial-vector normalizable wave functions subjects to vanishing UV and IR boundary conditions, they retain their *AdS* expressions except for the overall factor $1/kg_5^2$ which becomes $1/\hat{g}_5^2$.

THE LARGE- N BEHAVIOR OF THE HARD WALL MODEL

After having determined unambiguously the light quark mass and the chiral condensate entering the expression of the chiral symmetry breaking function (12) and clarified the

AdS/QCD dictionary as used in the *bottom-up* approach of the holographic models of QCD, let us focus on the large- N behavior of the Hard Wall model. A glance at the bulk fields, be it, for example, the normalizable vector mode $v_n(z)$ above or the vector bulk-to-boundary propagator for space-like momentum ($Q \equiv \sqrt{-q^2} > 0$):

$$\tilde{V}(Q^2, z) = Qz \left(K_1(Qz) + \frac{K_0(Qz_m)}{I_0(Qz_m)} I_1(Qz) \right) , \quad (18)$$

leads us to conclude that they do not scale with N . As a consequence, the square of the decay constants $F_{V_n}^2$, $F_{A_n}^2$ and f_π^2 in (17) scale as $O(N)$ which is reminiscent of what happens in Chiral Perturbation Theory with the large- N behavior implemented [6]. Let us go further by checking the large- N behavior of the electromagnetic $F_\pi(q^2)$ and gravitational $A_\pi(q^2)$ form factors of the pion [7] which take, in the chiral limit, very similar forms:

$$F_\pi(Q^2) = \frac{1}{kg_5^2} \frac{1}{f_\pi^2} \int_0^{z_m} dz z \tilde{V}(Q^2, z) \left[\left(\frac{\partial_z \Psi(z)}{z} \right)^2 + \frac{g_5^2 v(z)^2}{z^4} \Psi(z)^2 \right] , \quad (19)$$

$$A_\pi(Q^2) = \frac{1}{kg_5^2} \frac{1}{f_\pi^2} \int_0^{z_m} dz z \tilde{h}(Q^2, z) \left[\left(\frac{\partial_z \Psi(z)}{z} \right)^2 + \frac{g_5^2 v(z)^2}{z^4} \Psi(z)^2 \right] . \quad (20)$$

The so-called chiral field $\Psi(z) \equiv \phi_\chi(z) - \pi_\chi(z)$ is associated with the massless pseudo-scalar mode while $\tilde{h}(Q^2, z)$ is the tensor bulk-to-boundary propagator, analogous to (18), related to the fluctuations around the AdS_5 conformally flat metric (2). Besides their charge normalizations $F_\pi(0) = A_\pi(0) = 1$, the form factors do not scale with N which stems from the compensation mechanism between $1/kg_5^2 \sim O(N)$ and $1/f_\pi^2 \sim O(1/N)$. As for the *VPP* coupling constants, they follow writing the vector bulk-to-boundary propagator in terms of the vector mass poles m_{V_n} , of the residues F_{V_n} and of the normalizable eigenfunctions $v_n(z)$:

$$\tilde{V}(q^2, z) = -\sqrt{kg_5^2} \sum_{n=1}^{\infty} \frac{F_{V_n} v_n(z)}{q^2 - m_{V_n}^2 + i\epsilon} . \quad (21)$$

which does not scale, as it must be, with N . The *AdS* expression of the vector form factor and of the $g_{V_n \pi \pi}$ couplings follow:

$$F_\pi(q^2) = - \sum_{n=1}^{\infty} \frac{F_{V_n} g_{V_n \pi \pi}}{q^2 - m_n^2 + i\epsilon} \quad (22)$$

with

$$g_{V_n \pi \pi} = \frac{1}{\sqrt{kg_5^2}} \frac{1}{f_\pi^2} \int_0^{z_m} dz z v_n(z) \left[\left(\frac{\partial_z \Psi(z)}{z} \right)^2 + \frac{g_5^2 v(z)^2}{z^4} \Psi(z)^2 \right] . \quad (23)$$

As a result, the large- N counting rules are consistent in the *AdS/QCD* Hard Wall model as $g_{V_n \pi \pi} \sim O(1/\sqrt{N})$ such that we recover free, stable, non-interacting mesons at large N . Moreover, imposing that the pion decay constant coincides with the value

$f_\pi \simeq 92.4$ MeV in the chiral limit, one gets, with $1/z_m \simeq 323$ MeV, the quark condensate $\sigma \simeq (171 \text{ MeV})^3$. Because of the factor of k in the definition of $\bar{\sigma} = k\sigma$, the condensate turns out to be noticeably smaller than Hard Wall model estimates where this factor is usually omitted.

CONCLUSION

This talk aimed at clarifying some confusion regarding the large- N 't Hooft limit of the Hard Wall model of QCD which, used as a guideline, led us to study the different forms of the field/operator prescription within AdS/QCD . Such subtleties are not systematically taken into account when dealing with different 5d holographic models.

REFERENCES

1. J. M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, Adv. Theor. Math. Phys. **2** (1998) 231 [Int. J. Theor. Phys. **38** (1999) 1113] [arXiv:hep-th/9711200]; E. Witten, *Anti-de Sitter space and holography*, Adv. Theor. Math. Phys. **2** (1998) 253 [arXiv:hep-th/9802150]; S. S. Gubser, I. R. Klebanov & A. M. Polyakov, *Gauge theory correlators from non-critical string theory*, Phys. Lett. **B428** (1998) 105 [arXiv:hep-th/9802109].
2. J. Polchinski & M.J. Strassler, *Hard scattering and gauge/string duality*, Phys. Rev. Lett. **88** (2002) 031601 [arXiv:hep-th/0109174]. H. Boschi-Filho & N.R.F. Braga, *QCD/string holographic mapping and glueball mass spectrum*, Eur. Phys. J. **C32** (2004) 529 [arXiv:hep-th/0209080]; J. Erlich, E. Katz, D.T. Son & M.A. Stephanov, *QCD and a holographic model of hadrons*, Phys. Rev. Lett. **95** (2005) 261602 [arXiv:hep-ph/0501128]; L. Da Rold & A. Pomarol, *Chiral symmetry breaking from five dimensional spaces*, Nucl. Phys. **B721** (2005) 79 [arXiv:hep-ph/0501218].
3. A. Cherman, T.D. Cohen, E.S. Werbos, *The Chiral condensate in holographic models of QCD* [arXiv:hep-ph/0804.1096].
4. F. Jugeau, *Towards a consistent AdS/QCD dictionary* [arXiv:hep-ph/0902.3864].
5. P. Colangelo, F. De Fazio, F. Giannuzzi, F. Jugeau & S. Nicotri, *Light scalar mesons in the soft-wall model of AdS/QCD*, Phys. Rev. **D78** (2008) 055009 [arXiv:hep-ph/0807.1054].
6. A. Pich, *Colorless mesons in a polychromatic world*, published in "Tempe 2002, Phenomenology of large N_c QCD", 239-258, Ed. R. Lebed, World Scientific, Singapore, 2002 [arXiv:hep-ph/0205030].
7. H.R. Grigoryan & A.V. Radyushkin, *Pion form-factor in chiral limit of hard-wall AdS/QCD model*, Phys. Rev. **D76** (2007) 115007 [arXiv:hep-ph/0709.0500]; Z. Abidin & C.E. Carlson, *Gravitational Form Factors in the Axial Sector from an AdS/QCD Model*, Phys. Rev. **D77** (2008) 115021 [arXiv:hep-ph/0804.0214].